ELSEVIER

Developmental Brain Research 135 (2002) 105

DEVELOPMENTAL BRAIN RESEARCH

www.elsevier.com/locate/bres

Author Index

Avivi, C., Cui, S., Goren, S. and Goldstein, R.S.
Differences in neuronal differentiation between the transient cranial (Frorieps') and normal dorsal root ganglia (135) 19

Bothwell, M., see Karam, S.D. (135) 29
Boyd, A.W., see Karam, S.D. (135) 29
Brown, R.E., see Penner, M.R. (135) 71

Carrey, N., see Penner, M.R. (135) 71
Cottriall, C.L., see Vessey, K.A. (135) 79
Cui, S., see Avivi, C. (135) 19

de Rogalski Landrot, I., see Riviello, P. (135) 101
Dottori, M., see Karam, S.D. (135) 29

Fritzsch, B., see Maklad, A. (135) 1

Goldstein, R.S., see Avivi, C. (135) 19
Goren, S., see Avivi, C. (135) 19

Haber, S., see Lapidot, A. (135) 87
Henderson, J.T., see Karam, S.D. (135) 29
Holmes, G.L., see Riviello, P. (135) 101
Hongo, S., see Nakada, N. (135) 45
Hongo, S., see Ohki, T. (135) 55
Hutchinson, I., see Temple, E.C. (135) 65

Jinks, A.L., see Temple, E.C. (135) 65

Karam, S.D., Dottori, M., Ogawa, K., Henderson, J.T., Boyd, A.W., Pasquale, E.B. and Bothwell, M.
EphA4 is not required for Purkinje cell compartmentation (135) 29

Kuhn, H.-J., see Malz, C.R. (135) 39

Laing, D.G., see Temple, E.C. (135) 65
Lapidot, A. and Haber, S.
Effect of endogenous β-hydroxybutyrate on brain glucose metabolism in fetuses of diabetic rabbits, studied by ^{13}C magnetic resonance spectroscopy (135) 87

Maeda, A., see Nakada, N. (135) 45
Maeda, A., see Ohki, T. (135) 55
Maklad, A. and Fritzsch, B.
The developmental segregation of posterior crista and saccular vestibular fibers in mice: a carbocyanine tracer study using confocal microscopy (135) 1
Malz, C.R. and Kuhn, H.-J.
Calretinin and FMRFamide immunoreactivity in the nervus terminalis of prenatal tree shrews (*Tupaia belangeri*) (135) 39
McBrien, N.A., see Vessey, K.A. (135) 79
McFadyen, M.P., see Penner, M.R. (135) 71

Nakada, N., Hongo, S., Ohki, T., Maeda, A. and Takeda, M.
Molecular characterization of NDRG4/Bdm1 protein isoforms that are differentially regulated during rat brain development (135) 45
Nakada, N., see Ohki, T. (135) 55

Ogawa, K., see Karam, S.D. (135) 29
Ohki, T., Hongo, S., Nakada, N., Maeda, A. and Takeda, M.
Inhibition of neurite outgrowth by reduced level of NDRG4 protein in antisense transfected PC12 cells (135) 55
Ohki, T., see Nakada, N. (135) 45

Pasquale, E.B., see Karam, S.D. (135) 29
Penner, M.R., McFadyen, M.P., Pinaud, R., Carrey, N., Robertson, H.A. and Brown, R.E.
Age-related distribution of c-fos expression in the striatum of CD-1 mice after acute methylphenidate administration (135) 71
Pinaud, R., see Penner, M.R. (135) 71

Riviello, P., de Rogalski Landrot, I. and Holmes, G.L.
Lack of cell loss following recurrent neonatal seizures (135) 101
Robertson, H.A., see Penner, M.R. (135) 71

Takeda, M., see Nakada, N. (135) 45
Takeda, M., see Ohki, T. (135) 55
Temple, E.C., Hutchinson, I., Laing, D.G. and Jinks, A.L.
Taste development: differential growth rates of tongue regions in humans (135) 65

Vessey, K.A., Cottriall, C.L. and McBrien, N.A.
Muscarinic receptor protein expression in the ocular tissues of the chick during normal and myopic eye development (135) 79

0165-3806/02/$ – see front matter © 2002 Elsevier Science B.V. All rights reserved.
PII: S0165-3806(02)00406-6